

Supplemental Material

Antitubercular evaluation of root extract and isolated phytochemicals from *Lophira lanceolata* against two resistant strains of *Mycobacterium tuberculosis*

**Jeanne Louise Nkot^a, Dominique Serge Ngono Bikobo^{a,e*}, Auguste Abouem A
Zintchem^{a,b}, Norbert Mbabi Nyemeck II^{a,c}, Esther Del Florence Moni Ndedi^d, Patrick
Hervé Betote Diboué^d, Dieudonné Emmanuel Pegnyemb^a, Christian G. Bochet^e, and
Ulrich Koert^c**

^aDepartment of Organic Chemistry, Faculty of Science, University of Yaoundé I, P.O Box 812, Yaoundé, Cameroon;

^bDepartment of Chemistry, Higher Training College, University of Yaoundé I, P.O Box 47, Yaoundé, Cameroon;

^cPhilipps-Universität Marburg, Faculty of Chemistry, Hans-Meerwein-Strasse, D-35032 Marburg, Germany;

^dDepartment of Microbiology, Faculty of Science, University of Yaoundé I, P.O Box 812, Yaoundé, Cameroon;

^eDepartment Chemie, Universität Fribourg, CH du Musée 9, 1700 Fribourg, Switzerland

*Contact: Dominique Serge Ngono Bikobo

E-mail: ngonosh@gmail.com

Address: Department of Organic Chemistry, Faculty of Science, University of Yaoundé I, P.O Box 812, Yaoundé, Cameroon





Fig. S1 HR-ESI-MS spectrum of compound **1** (negative mode)




Fig. S2 HR-ESI-MS spectrum of compound **1** (Positive mode)




Fig. S3 ESI-MS spectrum of compound **1** (positive mode)




Fig. S4 ^1H -NMR spectrum of compound **1**




Fig. S5 ^{13}C -NMR spectrum of compound **1**